

A



A



				(795%)
1989	717			0.83 70.71, 0.97
2010	718			1.22 70.86, 1.73
2003	719			0.95 70.35, 2.60
1985	720			1.34 71.11, 1.62
1998	721			0.65 70.44, 0.95
2012	722			0.62 70.36, 1.08
2009	723			0.97 70.67, 1.40
2010	724			0.85 70.56, 1.29
2007	725			0.52 70.32, 0.85
2008	726			0.58 70.41, 0.82
2004	727			0.41 70.17, 1.02
2005	728			0.63 70.45, 0.89
2008	729			0.21 70.12, 0.37
2014	730			0.21 70.15, 0.29
2009	731			0.46 70.35, 0.60
2012	732			0.26 70.07, 0.97
2012	733			0.45 70.26, 0.78
2009	734			0.49 70.24, 1.00
2013	735			0.95 70.79, 1.14
2006	736			0.63 70.48, 0.82
2005	737			0.75 70.59, 0.95
2011	738	2010	0097	0.21 70.45, 0.63, 0.85

Handwashing with soap

Handwashing with soap is a key intervention to reduce the transmission of enteric pathogens. It is estimated that handwashing with soap can reduce the risk of diarrhoeal illness by 30-40% and the risk of acute respiratory infections by 15-25%. The effectiveness of handwashing with soap is dependent on the frequency and duration of handwashing, the use of soap, and the availability of clean water. Handwashing with soap is a simple and low-cost intervention that can be implemented in both urban and rural settings. It is a key component of a comprehensive WASH (Water, Sanitation, and Hygiene) program.

Excreta disposal

Excreta disposal is a key intervention to reduce the transmission of enteric pathogens. It is estimated that excreta disposal can reduce the risk of diarrhoeal illness by 10-20% and the risk of acute respiratory infections by 5-10%. The effectiveness of excreta disposal is dependent on the type of disposal method used, the frequency of disposal, and the availability of clean water. Excreta disposal is a simple and low-cost intervention that can be implemented in both urban and rural settings. It is a key component of a comprehensive WASH program.

Excreta disposal is a key intervention to reduce the transmission of enteric pathogens. It is estimated that excreta disposal can reduce the risk of diarrhoeal illness by 10-20% and the risk of acute respiratory infections by 5-10%. The effectiveness of excreta disposal is dependent on the type of disposal method used, the frequency of disposal, and the availability of clean water. Excreta disposal is a simple and low-cost intervention that can be implemented in both urban and rural settings. It is a key component of a comprehensive WASH program.




Fig. 2 Forest plot of risk ratios for the studies included in the meta-analysis




Fig. 3 Forest plot of risk ratios for the studies included in the meta-analysis

Availability of data and materials

Not applicable

30. 王德峰, F. 列奥蒂克, I. 康德. 王德峰讲读《康德》. 北京: 三联书店, 2014; 91(1):190-7.
31. 王德峰. 从《列奥蒂克》看《康德》. 哲学研究, 2009; 80(2):286-93.
32. 王德峰. 从《列奥蒂克》看《康德》. 哲学研究, 2012; 9(11):3806-23.
33. 王德峰. 从《列奥蒂克》看《康德》. 哲学研究, 1990; 0: 431.